

February 1, 2008

SISSA-ISAS 187/91/EP

Generalized Integrable Lattice Systems

C.S.XIONG

*International School for Advanced Studies
(SISSA/ISAS), Strada Costiera 11, I-34014 Trieste, Italy*

ABSTRACT

We generalize Toda-like integrable lattice systems to non-symmetric case. We show that they possess the bi-Hamiltonian structure.

1. INTRODUCTION

In the last two years there has been a great progress in understanding two dimensional quantum gravity coupled to minimal conformal models. One of the powerful tool to attack this problem is the so-called matrix model. It is well-known that one-matrix model is deeply connected with Toda-like (or Volterra-like) Lattice system[1,2,3], which is the simplest classical integrable model on the lattice[4,5]. Furthermore, on it one can realize a discretized Virasoro algebra as well as discretized KdV-equations, and probably it is related to a lattice version of Liouville theory[3,6].

These properties make it a natural question to consider generalizations of this model, maintaining the integrability, in particular, playing roles in the multi-matrix model and giving the possible formulation of the discrete W_n -algebra. A generalization has been proposed in[7], it is still an integrable symmetric lattice, but unfortunately it seems to have nothing to do with multi-matrix models. In this letter, we analyze a non-symmetric lattice and its associated discrete linear system. We prove that it is integrable. Its important role in multi-matrix models will be presented elsewhere[8].

2. NOTATIONS

Let us present here our notations. For any matrix M , we denotes its upper and lower triangular parts by $M_{\pm}$ respectively, and by $M_{(0)}$ means the diagonal line. We also introduce the definitions

$$M_{\theta} \equiv \frac{1}{2}M_{(0)} + M_{-}, \quad M_{\alpha} \equiv \frac{1}{2}M_{(0)} + M_{+}$$

$$M_a \equiv M_{(0)} + M_{+}, \quad I_{\pm} \equiv \sum_{i=0}^{\infty} E_{i,i\pm 1},$$

As usual $(E_{ij})_{kl} = \delta_{ik}\delta_{jl}$. We will consider throughout semi-infinte matrices, $0 \leq i, j < \infty$. Furthermore, for the later convenience, we introduce the natural

gradation

$$\deg[E_{ij}] \equiv j - i$$

$M_{[a,b]}$ denotes the diagonal lines of matrix M with degrees from “a” to “b”, while $M \in [a, b]$ means that the degrees of matrix M is only from “a” to “b”.

As we know, under matrix commutation, the semi-infinite matrices form an infinite dimensional algebra, we denote it by $gl_0(\infty)$, which has the following decomposition

$$gl_0(\infty) = \mathcal{N}_+ \oplus \mathcal{H} \oplus \mathcal{N}_-$$

the subalgebra $\mathcal{N}_+(\mathcal{N}_-)$ contains all the upper(lower) triangular matrices, while $\mathcal{H}$ represents the Abelian subalgebra formed by diagonal matrices. Furthermore, we introduce two subalgebras of $gl(\infty)$

$$\mathcal{P}_- \equiv \mathcal{H} \oplus \mathcal{N}_-, \quad \mathcal{P}_+ \equiv \mathcal{H} \oplus \mathcal{N}_+$$

3. THE SYMMETRIES OF THE DISCRETE LINEAR SYSTEMS

Let us begin with the following discrete linear system

$$\begin{cases} Q\xi = \lambda\xi \\ \frac{\partial}{\partial t_r}\xi = (Q^r)_\theta\xi \end{cases} \quad (1)$$

where the Jacobi matrix takes the following explicit form

$$Q_{ij} = \sqrt{R_j}E_{i,i+1}\delta_{j,i+1} + \sum_{l \geq 0}^n L_i^{(l)}E_{i,i-l}\delta_{j,i-l}, \quad i, j \geq 0 \quad (2)$$

Evidantly $Q \in [-n, 1]$ with arbitrary positive integer n , while R_i 's and $L_i^{(l)}$'s are coordinate variables of the system, whose equations of motion are

$$\frac{\partial}{\partial t_r}Q = [(Q^r)_\theta, Q] \quad \forall r \geq 1 \quad (3)$$

They are just the consistency conditions of the system (1), and we will call them the discrete KdV equations.

In this section we will consider two different types of the symmetries of the generalized discrete linear system: a gauge symmetry and a Weyl symmetry. The gauge symmetry preserves completely the dynamics of the system, while the Weyl scaling can be used to reduce the system (1) to the standard form.

Gauge Symmetry of The System

By gauge symmetry we mean the invariance of the system under the following transformations

$$\begin{cases} Q \longrightarrow \tilde{Q} = G^{-1}QG \\ \xi \longrightarrow \tilde{\xi} = G^{-1}\xi \end{cases} \quad (4)$$

where G is some invertible matrix such that the form of the system (1) remains unchanged, i.e.

$$\begin{cases} \tilde{Q}\tilde{\xi} = \lambda\tilde{\xi} \\ \frac{\partial}{\partial t_r}\tilde{\xi} = (\tilde{Q}^r)_\theta\tilde{\xi} \end{cases}$$

That is to say, the dynamics of the system does not change. In order to see what this symmetry is, we consider its infinitesimal form $G = 1 + \epsilon g$. The invariance requires that the matrix g should satisfy the equations

$$\begin{cases} \tilde{Q} = Q + \epsilon[Q, g], \\ \frac{\partial}{\partial t_r}g = [Q_\theta^r, g] - [Q^r, g]_\theta \end{cases} \quad (5)$$

Two solutions are

$$g_1 = \sum_k b_k Q^k, \quad g_2 = \sum_k c_k Q_\alpha^k, \quad \forall k \geq 1 \quad (6)$$

where b_k 's and c_k 's are time-independent constants. The effect of the transformations generated by g_1 's is only to rescale the basis by a λ -dependent factor, while the transformations generated by the g_2 's is in fact equivalent to tune the time parameters t_r 's. They form an infinite dimensional gauge group. We will see that, it is just this symmetry that leads to the integrability of the discrete linear system.

Weyl Scaling Symmetry

Another kind of symmetry of the system (1) is Weyl scaling invariance. Suppose that we choose a diagonal matrix G . After the transformation (4), the time evolution equations of ξ will transform to another form, but the equations of motion of the coordinates R_i 's and $L_i^{(l)}$'s keep unchanged. The only effect of this transformation is to rescale the basis of the vector ξ by a λ -independent factor. This symmetry has only one degree of freedom. Now let us choose the following special gauge $G_{ij} = \frac{1}{\sqrt{h_j}}\delta_{ij}$ (where we have introduced the quantities $h_j/h_{j-1} = R_j$, h_0 is an undetermined constant which is not important in our analysis), then the Jacobi matrix becomes

$$Q = I_+ + \sum_{l=0}^n A^{(l)}, \quad A^{(l)} \equiv \sum_{i=l}^{\infty} A_i^{(l)} E_{i,i-l} \quad (7a)$$

with

$$A_j^{(0)} = L_j^{(0)}, \quad A_j^{(l)} = L_j^{(l)} \sqrt{R_j R_{j-1} \dots R_{j-l+1}}, \quad \forall l \geq 1 \quad (7b)$$

In this gauge, it can be easily checked that the linear system (1) takes the following form

$$\begin{cases} Q\xi = \lambda\xi \\ \frac{\partial}{\partial t_r}\xi = (Q^r)_-\xi \end{cases} \quad (8)$$

The equations of motion (or consistency conditions of the system (8)) read

$$\frac{\partial}{\partial t_r} Q = [(Q^r)_-, Q] \quad \forall r \geq 1 \quad (9)$$

We see that the equations of motion of the coordinates R_j 's disappear, in fact it is implicitly involved in the equations of motion of the coordinates A_j 's. Both system (1) and system (8) appear in multi-matrix models[8].

4. INTEGRABILITY OF THE DISCRETE LINEAR SYSTEM (8)

As we know a dynamical system with n degrees of freedom is integrable if and only if there are n -independent conserved quantities in involution[4,5]. For a system with infinite many degrees of freedom, there should be infinite many independent conserved quantities in involution. One of the main approaches to show the integrability is the so-called bi-Hamiltonian method, i.e. one should prove that there exist at least two compatible Poisson brackets. Compatibility of two Poisson brackets means

$$\{H_{k+2}, f_Y\}_1(Q) = \{H_{k+1}, f_Y\}_2(Q) \quad (10)$$

for any Hamiltonian H_k . Hereafter we will follow this line to construct two compatible Poisson brackets, which gives the discrete KdV equations (9).

The First Poisson Bracket

Our strategy is as follows. We try to define such kind of Poisson structure, which gives the desired equations of motion, i.e. the discrete KdV-equations (9). In order to do this, let us define the trace operation on the matrix space*

$$Tr(M) \equiv \sum_{i=0}^{\infty} M_{ii}$$

which gives a natural inner scalar product on $gl_0(\infty)$. It is easy to see that the product is symmetric and invariant under the action of $gl_0(\infty)$

$$\begin{cases} Tr(AB) = Tr(BA) \equiv A(B) = B(A) \\ Tr(A[B, C]) = Tr([A, B]C) \end{cases}$$

with respect to this product, $\mathcal{P}_+$ and $\mathcal{P}_-$ are dual algebras, their elements being in 1-1 correspondence. On the other hand, the functions of $A_j^{(l)}$'s, which span a

* we suppose that this infinite summation is well-defined

functional space $\mathcal{F}$, can be defined as

$$f_X(Q) \equiv \text{Tr}(QX), \quad df_X(Q) = X \in \mathcal{P}_+$$

This is a function on the algebra $\mathcal{P}_-$. Consequently the algebraic structure of $\mathcal{P}_-$ determines the Poisson structure on $\mathcal{F}$ through the Konstant–Kirillov bracket

$$\{f_X, f_Y\}_1(Q) = Q([X, Y]), \quad X, Y \in \mathcal{P}_+ \quad (11)$$

Of course it is anti-symmetric and satisfies the Jacobi identity. Furthermore, one can show that with respect to this Poisson bracket, the conserved functions are

$$H_k = \frac{1}{k} \text{Tr}(Q^k) \quad dH_k = Q_{[0,n]}^{(k-1)} \quad \forall k \geq 1 \quad (12)$$

They are in involution, i.e.

$$\{H_{k+1}, H_{l+1}\}_1(Q) = 0$$

So they can be chosen as Hamiltonians, and give the desired equations of motion, i.e. the discrete KdV–Heirachies. On the other hand, these Hamiltonians generate the symmetry transformations on the functional space $\mathcal{F}$. In order to see its relation to the gauge transformations (6), we recall that the co-adjoint action of $\mathcal{P}_+$ on the functional space $\mathcal{F}$ is defined through its adjoint action on the coordinates space $\mathcal{P}_-$, i.e.

$$ad_Y^* f_X(Q) \equiv f_X(ad_Y(Q)) = X([Y, Q]) \quad X, Y \in \mathcal{P}_+$$

Obviously the symmetry on $\mathcal{F}$ generated by the Hamiltonians is nothing but the gauge symmetry on the coordinate space. Therefore, for a system with infinite degrees of freedom, there must exist an infinite dimensional gauge symmetry, such that we could find infinite many independent involutive conserved quantities.

The Second Poisson Bracket

In order to get the second Poisson bracket, our starting point is compatibility of two Poisson brackets. If we can represent the right hand side of eq.(10) in terms of dH_{k+1} , then we can extract the second Poisson bracket immediately. Consider the following matrix

$$\begin{aligned} F &= [Q, Q_a^{k+1}] \\ &= (Q(Q_a^k Q)_a - (Q Q_a^k)_a Q) + (Q Q_{[-1]}^k I_+ - I_+ Q_{[-1]}^k Q) \end{aligned}$$

From the equality

$$[Q, Q_a^k] = [Q_{-}^k, Q]$$

we obtain

$$Q_{[-1]}^k = \sum_{j=0}^{\infty} Q_{j+1,j}^k E_{j+1,j}, \quad Q_{j+1,j}^k = \sum_{l=0}^j [dH_{k+1}, Q]_{ll}$$

Therefore we get

$$\begin{aligned} \{H_{k+2}, f_Y\}_1(Q) &= Q([dH_{k+2}, Y]) = Y(F) \\ &= \langle (dH_{k+1} Q)_a Y Q \rangle - \langle (Q dH_{k+1})_a Q Y \rangle \\ &\quad + \sum_{i,j} Q_{ij} (Q_{j,j-1}^k - Q_{i+1,i}^k) Y_{ji} \end{aligned}$$

substituting dH_{k+1} by X , and introduce a X -dependent matrix

$$\mathcal{D} \equiv \sum_{j=0}^{\infty} \mathcal{D}_j E_{j+1,j} \quad \mathcal{D}_j \equiv \sum_{l=0}^j [X, Q]_{ll} \quad (13)$$

we finally obtain the second Poisson bracket

$$\begin{aligned} \{f_X, f_Y\}_2(Q) &= \langle (X Q)_a Y Q \rangle - \langle (Q X)_a Q Y \rangle \\ &\quad + \langle Q \mathcal{D} I_+ Y \rangle - \langle Q Y I_+ \mathcal{D} \rangle \end{aligned} \quad (14)$$

Writing explicitly in terms of the coordinates $A_j^{(l)}$'s, we have

$$\begin{aligned}
\{A_i^{(a)}, A_j^{(b)}\}_2 &= A_j^{(a+b+1)} \delta_{i,j-b-1} - A_i^{(a+b+1)} \delta_{i,j+a+1} \\
&\quad + \sum_{l=0}^a (A_i^{(l)} A_j^{(a+b-l)} \delta_{i,j-b+l} - A_i^{(a+b-l)} A_j^{(l)} \delta_{i,j+a-l}) \\
&\quad + A_i^{(a)} A_j^{(b)} \sum_{l=j-b}^j (\delta_{il} - \delta_{i-a,l})
\end{aligned} \tag{15}$$

here we understand $A_i^{(r)} = 0$, if $r > n$. It is tiresome but straightforward to prove that the above Poisson bracket is antisymmetric and satisfies the Jacobi identity. The following is a simple example for $n = 2$ and

$$Q_{ij} = \delta_{j,i+1} + S_j \delta_{ij} + L_j \delta_{j,i-1} + A_j \delta_{j,i-2}$$

Eq.(15) gives us the following Poisson brackets

$$\{S_i, S_j\}_2 = L_j \delta_{i,j-1} - L_i \delta_{i,j+1}, \tag{16a}$$

$$\{S_i, L_j\}_2 = S_i L_j (\delta_{i,j-1} - \delta_{ij}) + A_j \delta_{i,j-2} - A_i \delta_{i,j+1}, \tag{16b}$$

$$\{S_i, A_j\}_2 = S_i A_j (\delta_{i,j-2} - \delta_{ij}), \tag{16c}$$

$$\{L_i, S_j\}_2 = L_i S_j (\delta_{ij} - \delta_{i,j+1}) + A_j \delta_{i,j-1} - A_i \delta_{i,j+2}, \tag{16d}$$

$$\{L_i, L_j\}_2 = L_i L_j (\delta_{i,j-1} - \delta_{i,j+1}) + S_i A_j \delta_{i,j-1} - A_i S_j \delta_{i,j+1}, \tag{16e}$$

$$\{L_i, A_j\}_2 = L_i A_j (\delta_{i,j-2} + \delta_{i,j-1} - \delta_{ij} - \delta_{i,j+1}), \tag{16f}$$

$$\{A_i, S_j\}_2 = A_i S_j (\delta_{ij} - \delta_{i,j+2}), \tag{16g}$$

$$\{A_i, L_j\}_2 = A_i L_j (\delta_{i,j-1} + \delta_{ij} - \delta_{i,j+1} - \delta_{i,j+2}), \tag{16h}$$

$$\{A_i, A_j\}_2 = A_i A_j (\delta_{i,j-2} + \delta_{i,j-1} - \delta_{i,j+1} - \delta_{i,j-2}), \tag{16i}$$

From our construction of the Poisson brackets, at the first glance it seems to be possible to deduce higher order Poisson brackets. However it is very difficult to work it out, and, furthermore, the deduced higher order Poisson brackets probably

do not satisfy Jacobi identity. The only exception is the ansatz $n = 1, r = 2$, in this case, we denote

$$Q = I_+ + S + R$$

Then playing the same game as before, we can get the third Poisson bracket of the Toda-like lattice

$$\begin{aligned} \{f_X, f_Y\}_3(Q) = & \frac{1}{2}(\langle (XQ^2)_a YQ \rangle - \langle (Q^2 X)_a QY \rangle \\ & + \langle (QXQ)_a YQ \rangle - \langle (QXQ)_a QY \rangle \\ & \langle YQC_1 \rangle - \langle QYC_2 \rangle) \end{aligned} \quad (17)$$

where

$$\begin{aligned} B &\equiv \sum_{j=0}^{\infty} B_j E_{j+2,j}, & B_j &\equiv \sum_{l=0}^j [X, Q_a]_{l+1,l} \\ \mathcal{D} &\equiv \sum_{j=0}^{\infty} \mathcal{D}_j E_{j+1,j}, & \mathcal{D}_j &\equiv \sum_{l=0}^j [X, Q]_{ll} \end{aligned}$$

and

$$\begin{aligned} C_1 &\equiv BI_+^2 + I_+BI_+ + \mathcal{D}Q_+^2 2I_+ \mathcal{D}I_+ + I_+ \mathcal{D}S + S \mathcal{D}I_+ \\ C_2 &\equiv I_+^2 B + I_+BI_+ + Q_+^2 \mathcal{D} + 2I_+ \mathcal{D}I_+ + I_+ \mathcal{D}S + S \mathcal{D}I_+ \end{aligned}$$

The Poisson brackets of the coordinates are as follows

$$\begin{aligned} \{R_i, R_j\}_1 &= 0, & \{S_i, S_j\}_2 &= 0. \\ \{R_i, S_j\}_1 &= R_i(\delta_{j,i+1} - \delta_{i,j}) \end{aligned} \quad (18)$$

and

$$\{R_i, R_j\}_2 = R_i R_j (\delta_{j,i+1} - \delta_{i,j+1}) \quad (19a)$$

$$\{R_i, S_j\}_2 = R_i S_j (\delta_{j,i+1} - \delta_{i,j}) \quad (19b)$$

$$\{S_i, S_j\}_2 = R_i \delta_{j,i+1} - R_j \delta_{i,j+1}. \quad (19c)$$

as well as

$$\{R_i, R_j\}_3 = 2R_i R_j (S_i \delta_{i,j-1} - S_j \delta_{i,j+1}) \quad (20a)$$

$$\begin{aligned} \{R_i, S_j\}_3 &= R_i R_j (\delta_{i,j-1} + \delta_{i,j}) - R_i R_{j+1} (\delta_{i,j+1} + \delta_{i,j+2}) \\ &\quad + R_i S_j^2 (\delta_{i,j} - \delta_{i,j+1}) \end{aligned} \quad (20b)$$

$$\{S_i, S_j\}_3 = (S_i + S_j)(R_j \delta_{i,j-1} - R_i \delta_{i,j+1}) \quad (20c)$$

These Poisson brackets have been derived in [3] by a straightforward calculation, while now they have been obtained from the sytematic analysis. In the continuum limit, if we set the infinitesimal parameter $\epsilon = \frac{1}{N}$, and suppose that

$$\begin{cases} R_j \longrightarrow 1 + \frac{1}{2}\epsilon^2(u(x) + v(x)) \\ S_j \longrightarrow 2 + \frac{1}{2}\epsilon^2(u(x) - v(x)) \end{cases} \quad (3.27)$$

after introducing a new Poisson bracket

$$\{, \} \equiv \frac{1}{4} \{, \}_3 - \{, \}_2$$

we get two copies of Virasoro algebras

$$\{u(x), u(y)\} = \frac{1}{2}(\partial^3 + 4u(x)\partial + 2u'(x))\delta(x - y) \quad (3.28a)$$

$$\{v(x), v(y)\} = -\frac{1}{2}(\partial^3 + 4v(x)\partial + 2v'(x))\delta(x - y) \quad (3.28b)$$

$$\{u(x), v(y)\} = 0. \quad (3.28c)$$

5. INTEGRABILITY OF THE SYSTEM (1)

In the previous section, we have gauged away the Weyl symmetry keeping in this way one degree of freedom. So the coordinates R_j 's do not appear in the Poisson brackets as well as in the discrete KdV equations. Now let us see what is the suitable Poisson brackets including them explicitly.

If we do not fix the Weyl symmetry, then we have the discrete linear system (1), whose integrability can be analysed in almost the same way as before. The only difference is that in this case $Q \in [-n, 1]$ but the tangent vector belongs to $[-1, n]$, so we should make use of R-matrix of $gl_0(\infty)$, which acts in the following way

$$\begin{cases} \mathcal{R}(E_{ij}) = E_{ij}, & i < j \\ \mathcal{R}(E_{ii}) = 0 \\ \mathcal{R}(E_{ij}) = -E_{ij}, & i > j. \end{cases} \quad (21)$$

which defines the following Lie-algebraic structure on the coordinate space

$$[X, Y]_{\mathcal{R}} = [\mathcal{R}(X), Y] + [X, \mathcal{R}(Y)]$$

Now let us consider the induced symmetry transformation on the functional space by gauge symmetry (6), which is in the coordinate space.

$$f_X(Q) \longrightarrow G^{-1} f_X(Q) G \equiv f_X(G^{-1} Q G) \in \mathcal{F}$$

In particular, set $g = Q_{\alpha}^k$, and introduce the quantities

$$H_k \equiv \frac{1}{k} \text{Tr}(Q^k), \quad \forall k > 0$$

then

$$\delta f_X(Q) = \epsilon X([Q, g]) = \epsilon \frac{1}{2} Q([dH_{k+1}, X]_{\mathcal{R}})$$

which can be considered as the transformation generated by function H_{k+1} through

some Poisson bracket, which is the first Poisson bracket

$$\{f_X, f_Y\}_1(Q) = \frac{1}{2}Q([X, Y]_{\mathcal{R}}) \quad (22)$$

with respect to this Poisson bracket, H_k 's are the conserved quantities. Choosing one of them as Hamiltonian we recover the KdV–Heirachioies. Obviously, this Poisson bracket is antisymmetric and satisfies the Jacobi identity.

In order to construct the second Poisson bracket, once again we use compatibility condition (10)(since it is valid for any integrable system). Finally the straightforward computation shows that the second Poisson bracket is

$$\{f_X, f_Y\}_2(Q) = \frac{1}{2} < QX\mathcal{R}(QY) - \mathcal{R}(YQ)XQ > \quad (23)$$

Now let us consider once again the $n=2$ case

$$Q_{ij} = \sqrt{R_j}\delta_{j,i+1} + S_j\delta_{ij} + l_j\delta_{j,i-1} + a_j\delta_{j,i-2}$$

where

$$\sqrt{R_j}l_j = L_j, \quad \sqrt{R_j R_{j-1}}a_j = A_j$$

then the Poisson bracket (23) gives the Poisson algebra which contains (16) as a sub-algebra and the following ones

$$\{S_i, R_j\}_2 = S_i R_j (\delta_{i,j-1} - \delta_{ij}), \quad (24a)$$

$$\{L_i, R_j\}_2 = L_i R_j (\delta_{i,j-1} - \delta_{i,j+1}), \quad (24b)$$

$$\{A_i, R_j\}_2 = A_i R_j (\delta_{i,j-1} - \delta_{i,j+2}), \quad (24c)$$

$$\{R_i, R_j\}_2 = 0, \quad (24d)$$

There are a few unclear problems. At first, it is interesting to investigate the relation between the continuum version of the above Poisson brackets and the

Gelfand–Dickii Poisson brackets. Secondly these non-symmetric lattices probably give a realization of discrete W_n -algebras. One may also ask if there exist some possible reduction of this model, which leaves only one degree of freedom in $Q_{[-n,1]}, \forall n \geq 2$. This reduction might be important in multi-matrix models[8].

Acknowledgements

I would like to thank Prof. L.Bonora for valuable suggestions and stimulating discussions. I also wish to thank Prof.M.Martellini for discussions and comments.

References

- [1] E.J.Martinec, “On the Origin of Integrability of Matrix Models”, Chicago preprint, EFI-90-67
- L.Alvarez-Gaume, “The Integrability in Random Matrix Models”, CERN-TH.6123/91
- [2] A.Gerasimov, A.Marshakov, A.Mironov, and A.Orlov, “Matrix Models of Two Dimensional Gravity and Toda Theory”, ITEP preprint (1990)
- [3] L.Bonora, M.Martellini and C.S.Xiong, SISSA preprint EP-107/91
- [4] L.D.Faddeev, L.A.Takhtadzyan, “Hamiltonian Methods in the Theory of Solitons”, Springer(1987)
- [5] O.Babelon and C.Viallet, Lecture Notes in SISSA (1989)
- [6] O.Babelon and L.Bonora, Phys.Lett.**253B**(1991)365
- [7] J.Avan, Brown preprint BROWN-HET-812
- [8] C.S.Xiong, SISSA preprint